

Available Procedures At Manchester Private Hospital

BREAST SURGERY

- Breast Enlargement
- Breast Reduction
- Breast Uplift
- Fat Transfer Breast Aug
- Breast Re-Augmentation
- Breast Asymmetry
- Breast Reconstruction
- Transgender Breast Surgery
- Nipple Correction

BODY SURGERY

- Liposuction
- Vaser Liposuction
- Vaser Hi-def
- Tummy Tuck
- Mini Tummy Tuck
- Arm Lift
- Thigh Lift
- 360 Liposuction
- Body Lift Surgery
- Buffalo Hump Removal
- Fat Transfer
- Lipoedema Liposuction
- Pubic Fat Pad Removal
- Gynecomastia
- J Plasma Renuvion
- Vaginal Tightening
- Labiaplasty
- Varicose Veins
- Pelvic Floor Repair
- Brazilian Butt Lift
- Mummy Makeover

FACE SURGERY

- Facelift
- Facial Lipo
- Chin Implants
- Eyebrow Lift
- Nose Surgery
- Eyebag Removal
- Ear Correction
- Septorhinoplasty
- Neck Lift Surgery



MANCHESTER
PRIVATE HOSPITAL

Post-Operative Advice Blepharoplasty

WHEN IS MY MEDICATION NEXT DUE?

Pain Relief - Time _____


Antibiotic - Time _____

Excellent   Trustpilot

CONTACT US

Manchester Private Hospital

New Court, Regents Place, Windsor Street Salford,
Greater Manchester, M5 4HB.

 0161 507 8822

 manchesterprivatehospital.uk

manchesterprivatehospital.uk

For any Post Operative Queries

0161 507 8822

FOLLOWING YOUR BLEPHAROPLASTY SURGERY

postoperative bruising and swelling is normal and should subside over the next 3-6 weeks.

You may leave the hospital with steri-strips (small paper thin) dressings over your eye lid incisions. Please leave these intact until you come back to see the nurse from 7 days after your surgery.

Do not remove these as the stitch may be attached to them and removal may pull the stitch out.

It is normal for your vision to be blurred for a few days following surgery therefore it is advisable to rest at home for 7 days post op. Please do not drive during this time.

Your eyes will be light sensitive also, so please wear dark sunglasses in day light for the first few weeks following surgery and be aware that new scars will burn easily in sunlight.

Avoid heavy lifting and bending down without bending your knees as this can cause bleeding due to increased pressure.

Avoid strenuous exercise for at least 6 weeks after surgery as this can increase risk of bleeding and poor healing.

You may need around 2 weeks off work depending on your job role. Make up can be worn after 2-3 weeks. Do not drive your car or drink alcohol for at least 1 week after surgery, the pain relief may 'slow down' your normal reactive response.

Please sleep in a semi-upright position to aid reduction in swelling overnight.

A cold compress can be gently applied over the face and eyes following surgery.

You may wish to buy facial/eye cold packs these are available from Boots the pharmacy and are very soothing following surgery. They also help reduce swelling in the first 24 hours post operatively. Do not apply frozen cool packs directly to skin as this can harm the skin, first wrap the cool pack with a clean cotton covering or fresh gauze.

You will be given pain relief prescribed by the surgeon.

Take these when required. Usually paracetamol and codeine may be prescribed but after 2-3 days if you no longer require them stop. Medication containing codeine can cause you to feel low in mood and can cause constipation so when you no longer need them stop taking them.

To help prevent constipation after surgery eat a good diet with fruit and vegetables. Drink plenty of fluids and walk gently at least hourly throughout your day. Staying mobile is essential in reducing risk of deep vein thrombosis following surgery.

It is important to wear your anti thrombosis stockings until you are fully mobile, minimum 2-3 days following surgery and longer if you are inactive.

Signs of a blood clot called deep vein thrombosis (DVT) is calf pain and swelling. If this happens you must attend Accident and Emergency for investigations.

Deep vein thrombosis or DVT can lead to a blood clot in the lung called PE or Pulmonary Embolism which is life threatening.

The signs of PE are chest pain and shortness of breath. If you experience these symptoms, you must call 999 as this is a medical emergency.

You may be given an antibiotic cream called Chloramphenicol to apply to your eye lids and eye line twice daily am and pm.

This is very soothing and helps reduce risk of infection however it can blur your vision as it is an ointment.

You may also be given oral antibiotics please complete the prescribed course.

Please inform us of any negative side effects, worries or concerns that you experience.

You may shower the following day after surgery but please keep the eye lids dry and leave steri-strips in place.

Blinking tightly can help reduce swelling by squeezing the fluid out of the eye lids. You can try this morning and night.

Bruising and swelling is normal and is to be expected, this should subside within 4-6 weeks after surgery. The cosmetic appearance will improve as swelling settles and can take around 3-6 months to see the final/optimum result when scarring has settled.

If your eyes become sticky, blurred or infected please let us know immediately so we can make an appointment for you to see the surgeon.

Your eyes may not fully close to begin with due to swelling and tightness of stiches. If they become dry overnight, you may apply artificial tears from the pharmacy to help

keep eyes lubricated. This will settle over time.

DO NOT SMOKE OR BE AROUND SMOKERS FOR AT LEAST 3 MONTHS AFTER SURGERY AS THIS CAN LEAD TO INFECTION AND POOR HEALING.

<p>EMERGENCY</p> <p>IN CASE OF EMERGENCY PLEASE DIAL 999 AND ATTEND ACCIDENT AND EMERGENCY THIS INCLUDES SUDDEN HIGH TEMPERATURE, BLEEDING, CHEST PAIN AND BREATHLESSNESS.</p>
--